
AD - A 13	2 410	EASTERN- POLAR OC	WESTERN ARCT	IC SEA ENTER W	ICE ANA	LYSIS	1982 (U) 1982	NAVAL	. 1	14	1
UNCLAS	SIFIED						F/0	8/12	NL		
		3 /4				7) () ()	75.7		·	15.1	2.1
777		775	*> > ?	- 15g c	უი. ა	47) r	"> ·	-Æ} ⁻	"> <u>*</u>	- 87 c	r) A
-5-5-		177) (*	- #5	27.0°	<u>ත</u> ුරු	- 27) /	***	77	ウ ト	57	ijy ir
وتو ، وت	×	ودي	19919	87) / .	30	<i>5</i> 77	32	J.;		Z):	77.7
79 (7)	*		191175	. 51	30°.	451	У ^л .				
			A .		e. V		1.		,	\ <u>.</u>	
			3 3	7,	•	ь	** ' >	. *, *	Ta Fa	*	*. V.



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS - 1963 - A

UNCLASSIFIED

EASTERN—WESTERN
ARCTIC SEA ICE ANALYSIS
1982

PREPARED BY NAVAL POLAR OCEANOGRAPHY CENTER, SUITLAND, MD

COPY


3

AD A 1

PREPARED FOR COMMANDER,

NAVAL OCEANOGRAPHY COMMAND NSTL STATION, BAY ST. LOUIS, MS 39529 DISTRIB Approv

88 08



STERN E ANALYSIS

APHY CENTER,

DTIC ELECTE SEP 13 1983

DISTRIBUTION STATEMENT A

Approved for public releases
Distribution Unlimited

OMMAND DUIS, MS 39529

83 08 10 035

The U.S. Navy has a long and colorful history of polar exploration and currently is an active participant in the growing national activity in the Arctic and Antarctic. The strategic importance and increased demand for the natural resources of these areas have resulted in a greater requirement for environmental information. In 1976, the National Oceanic and Atmospheric Administration (NOAA) joined the Navy in forming a Joint Ice Center (JIC) to combine efforts and resources in sea ice analysis and forecasting.

Through 1972, reliable sea ice information in the polar regions was based on a relatively few shore station and ship reports augmented by limited aerial reconnaissance data. These data were further restricted mainly to the relatively small areas observed primarily during the ship operating seasons. The advent of high resolution satellite imagery combined with the ground truth of conventional observations has in recent years enabled a description of the polar ice fields on a semi-synoptic scale in both polar regions.

This publication is the ninth in a continuing yearly series of Arctic sea ice atlases prepared in the Joint Ice Center at the Naval Polar Oceanography Center, Suitland. The atlas contains weekly charts depicting Northern Hemisphere and Great Lakes ice conditions and extents. The information presented was prepared under operational time constraints principally from satellite imagery supplemented by conventional observations. The Great Lakes ice analyses are derived principally from information provided by the National Weather Service Forecast Office, Ann Arbor, Michigan. Table 1 located on the inside back cover summarizes satellite data availability for 1982.

The purpose of with reliable wee limitations and oth various forms of procedures:

(1) a. Conventiona naissance observation timeliness;

the most recent and conventional with sa

(3) e Where suffilimit, an estimated availability the icice drift data and no data availability

	Just	ifice	tion	
	By T	er	叶	on f
		ribut		
	Ava	ilabi	11t3	Codes
		Ava	il a	nd/or
\	Dist	S	peci	al
)	n			

Accession For

NTIS GRALI

DTIC TAB Unannounced



FOREWORD

history of polar t in the growing . The strategic escurces of these for environmental tmospheric Adminlce Center (JIC) and forecasting.

the polar regions reports augmented ata were further observed primarily f high resolution h of conventional ption of the polar gions.

g yearly series of enter at the Naval tains weekly charts ce conditions and er operational time supplemented by alyses are derived hal Weather Service ated on the inside or 1982, The purpose of this atlas is to provide operators and researchers with reliable weekly hemispheric ice analyses. Satellite data limitations and other difficulties involved in manually synthesizing various forms of ice data have evolved the following analysis procedures:

(1) a. Conventional shore station, ship and aerial ice reconnaissance observations are plotted on base charts and evaluated for timeliness;


Satellite data is analyzed for ice information content, with the most recent and highest resolution considered first. Synthesis of conventional with satellite data yields the final analysis;


(3) e Where sufficient data is not available to define the sea ice limit, an estimated boundary is depicted. After one week of no data availability the ice edge position is based on analyzed theoretical ice drift data and other diagnostic aids. During subsequent weeks of no data availability the ice edges approach seasonal mean positions.

Acces	ion For				
ntis	GRALI				
DTIC :	PAB	<u>"</u> "			
	ounced				
Justi	fication	L			
By Per Ltr. on file					
Distribution/					
Avai	lability	Codes			
Avail and/or					
Dist	Speci	al			
1					

Joint Ice Center Naval Polar Oceanography Center 4301 Suitland Road Washington, DC 20390











)


-































2_




































L




__











_

























































j





-





































/























_


384








-










































_











) }














UNCLASSIFIED				F/G 8/12 NL		
	15 15 15 K	17/7/2	.5.	13 PO P	500	
i e e	14 84	ر على عا	4 7 7	477	4 7 7	i ni n
i mi m	477	4 7 7	4 7	1,7,7	1, 7, 7	1, 78, 79
1 × ×	1 7 7	1 7 7	4 7 7	1 7 7	(7)	1,7,7
4.87	A N P	1 7	4 4 9	14. 7 P	476	3.
;	1 1 1					





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS - 1963 - A




























-















-














*















































































































































































!




























































AD-A132 4	P	ASTERN OLAR O	I - WESTERI ICE ANDGR	N ARCI	IC SEA	ICE AN	ALYSIS TON DC	1982 (U 1982) NAVAL	3	14		4
UNCLASSIF	IED							F/	G 8/12	NL			
		7*		¥**		7 74	•	. 97		מל		, pp	
(4 , 5	35.7	7		⊘	30	7		END Are notes 9 83	CONT				-
'									•				
						_							





MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS - 1963 - A
























-


34
































TABLE I SATELLITE DATA UTILIZED DURING 1982

Time P	eri od	Satell	ite Remote Se	Resolution	6		
From	То	Sensor Platform	Sensor Type	Spectral Region	Resolution	Coverage	
1-82	1282	NOAA-6	AVHRR HRPT/LAC VIS NIR IR IR	0.55-0.68 um 0.73-1.10 um 3.55-3.93 um 10.5-11.5 um	1 km	Regional	
			GAC VIS IR	0.55-0.68 um 10.5-11.5 um	4km	Global	
1-82	12-82	NOAA-7	AVHRR HRPT/LAC V1S NIR IR IR	0.58-0.68 um 0.73-1.10 um 3.55-3.93 um 10.3-11.3 um	11km	Regional	
			GAC VIS IR	0.58-0.68 um 10.3-11.3 um	4km	Global	
1-82	11-82	NIMBUS-5 *	ESMR	1.55 cm	25km	Global	
11-82	12-82					Regional	
1-82	12-82	NIMBUS-7	SMMR	0.81 cm 1.66 cm	50km	Global	

Abbreviations and Acronyms

AVHRR - Advanced Very High Resolution Radiometer

cm - Centimeter
ESMR - Electrically Scanning Microwave Radiometer
GAC - Global Area Coverage


IRPT - High Resolution Picture Transmission
IR - Infrared
lum - Kilometer

LAC - Local Area Coverage
NIR - Near Infrared
SMAR - Scanning Multifrequency Microwave Radiometer

VHRR - Very High Resolution Radiometer

VIS - Visual um - Micrometer


^{*} NIMBUS-5 recorders failed on 23 November 1982, limiting data to the Alaska region.



AD-A132 410 EASTERN-WESTERN ARCTIC SEA ICE ANALYSIS 1982(U) NAVAL POLAR OCEANOGRAPHY CENTER WASHINGTON DC MAR 83

UNCL'ASSIFIED

F/G 8/12 NL



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS - 1963 - A

SUPPLEMENTAR

INFORMATION

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2. GOVT ACCESSION NO.	
4. TITLE (end Subtitle)	5. TYPE OF REPORT & PERIOD COVERED
Eastern-Western Arctic Sea Ice Analysis 1982	Reference Report 1982
1302	6. PERFORMING ORG, REPORT NUMBER
7. AUTHOR(*) Naval Polar Oceanography Center, Suitland, MD	8. CONTRACT OR GRANT NUMBER(*)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Oceanography Command Detachment Federal Bldg Asheville, NC 28801	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE Mar 83
Commander Naval Oceanography Command NSTL. MS 39529	13. NUMBER OF PAGES
14. MONITORING AGENCY NAME & ADDRESS(II dillerent from Controlling Office)	15. SECURITY CLASS. (of this report)
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)	

Approved for public release; distribution unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

18. SUPPLEMENTARY NOTES

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

Sea ice, polar ice fields, satellite imagery, concentration, stage of development, fast ice, concentration of thickness, theoretical thickness, Arctic

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

These are approximately 7-days analysis of sea ice prepared by the Naval Polar Oceanography Center, Suitland, MD. Included are ice concentration and ice thickness (age).

DD 1 JAN 73 1473

EDITION OF I NOV 65 IS OBSOLETE

5 N 0102- LF- 014- 6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered